

Raise the bed fence.





The moment when you look away from your child!

Even for a brief moment...

During hospitalization, children are more anxious than usual due to the different environment and physical condition, and they behave differently from usual. Is the bed fence raised up to the top?



Is there anything on the bed that your child can step on and climb out of bed? Is there anything that may cause your child to choke or suffocate? Please check again for safety.

Hold hands when walking!!

You may have to change diapers more often than usual

when an IV straw is

connected to you.

You may carelessly forget that an IV straw is connected to you.

Children may forget about the IV line and start running when they find something interesting or feel anxious.





They may feel unsteady on their feet unexpectedly when walking after staying in bed for a long period of time. With this in mind, please hold hands with your child for safety.

Wear shoes that are comfortable to walk in.



come off easily?

Can your shoes cause you to trip and fall?

Your child may not be able to walk like they usually do after being in bed for a long period of time. If your child wears shoes that come off easily, he/she may trip and fall. Even if your

Be more careful and stay safe!



child did not have any problems with walking at home, it is best to check again for safety. sandals X slippers X indoor shoes O sneakers with Velcro O



- 制作日 2023.1.10
- 制作 女子美術大学 アート・デザイン表現学科 ヒーリング表現領域・アートプロデュース表現領域3年 野呂千花子 日渡咲絢 福井さやか 藤田睦美 山崎友花
- 翻訳協力 国立国際医療研究センター病院 国際診療部
- 文 国立国際医療研究センター病院 小児科

©National Center for Global Health and Medicine. All rights reserved.